

University of South Carolina Scholar Commons

Faculty Publications

Chemistry and Biochemistry, Department of

3-2007

Poly[$\text{diaqua}(\mu_4\text{-3-fluorophthalato-}\kappa^4\text{O:O:O':O'')cadmium(II)}$]

Diana Rishmawi
Francis Marion University

Katie Lewis
Francis Marion University

Mark D. Smith
University of South Carolina - Columbia

LeRoy Peterson Jr.
Francis Marion University

Hans Conrad zur Loye
University of South Carolina - Columbia, zurloye@sc.edu

Follow this and additional works at: https://scholarcommons.sc.edu/chem_facpub



Part of the [Chemistry Commons](#)

Publication Info

Published in *Acta Crystallographica Section E*, Volume E63, Issue 3, 2007, pages m695-m697.

Copyright © International Union of Crystallography

DOI: 10.1107/S1600536807004898

Publisher's Version: <http://dx.doi.org/10.1107/S1600536807004898>

This Article is brought to you by the Chemistry and Biochemistry, Department of at Scholar Commons. It has been accepted for inclusion in Faculty Publications by an authorized administrator of Scholar Commons. For more information, please contact dillarda@mailbox.sc.edu.

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

Editors: **W. Clegg** and **D. G. Watson**

Poly[*diaqua*(μ_4 -3-fluorophthalato- $\kappa^4 O:O:O':O'$)cadmium(II)]

Diana Rishmawi, Katie Lewis, Mark D. Smith, LeRoy Peterson Jr and Hans-Conrad zur Loye

Copyright © International Union of Crystallography

Author(s) of this paper may load this reprint on their own web site provided that this cover page is retained. Republication of this article or its storage in electronic databases or the like is not permitted without prior permission in writing from the IUCr.

Poly[diaqua(μ_4 -3-fluorophthalato- κ^4 O:O:O':O'')-cadmium(II)]

Diana Rishmawi,^a Katie Lewis,^a
Mark D. Smith,^b LeRoy Peterson
Jr^a and Hans-Conrad zur Loye^{b*}

^aChemistry Department, Francis Marion University, Florence, South Carolina 29501, USA, and ^bDepartment of Chemistry and Biochemistry, University of South Carolina, Columbia, South Carolina 29208, USA

Correspondence e-mail: lpeterson@fmarion.edu

Received 15 December 2006

Accepted 30 January 2007

Key indicators

Single-crystal X-ray study

$T = 294$ K

Mean $\sigma(\text{C}-\text{C}) = 0.003$ Å

R factor = 0.017

wR factor = 0.042


Data-to-parameter ratio = 12.0

For details of how these key indicators were automatically derived from the article, see <http://journals.iucr.org/e>.

The crystal structure of the title compound, $[\text{Cd}(\text{C}_8\text{H}_3\text{FO}_4)(\text{H}_2\text{O})_2]_n$, consists of polymeric sheets formed by the bridging of octahedrally coordinated Cd^{II} by carboxylate O atoms of the 3-fluorophthalate (3-fpt^{2-}) ligand. The layers exhibit hydrogen bonding between each of two coordinated water molecules and two O atoms of the ligand. Adjacent sheets are connected through π - π interactions.

Comment

Polycarboxylates represent a versatile class of ligands for the construction of metal-organic coordination polymers (Ye *et al.*, 2005). Their versatility stems from the variety of bridging modes of the carboxylate group (Mehrotra & Bohra, 1983). Many of these solids exhibit properties such as gas absorption, catalytic activity, and luminescence (Rowell *et al.*, 2004; Wasuke *et al.*, 2005; Kim *et al.*, 2004). With this in mind, the title compound, (I), was prepared as a part of our ongoing efforts to construct new coordination polymers employing polycarboxylate ligands.



The asymmetric unit consists of one Cd^{II} ion, a 3-fluorophthalate (3-fpt^{2-}) dianion, and two coordinated water molecules. The Cd^{II} ion exhibits an all-oxygen coordination in a distorted octahedral environment (Fig. 1). The axial sites are defined by a water O atom and a 3-fpt^{2-} O atom, while the equatorial sites are defined by a water O atom and three carboxylate O atoms from three equivalent 3-fpt^{2-} ligands.

The Cd—O bond distances are normal (Table 1), with an average value of 2.3031 (18) Å. All other distances and angles are comparable with those in a similar Cd^{II} -phthalate coordination polymer (Vaz *et al.*, 1996).

In (I) there are $\text{O}-\text{H}\cdots\text{O}$ hydrogen bonds involving three H atoms of the two water molecules and two O acceptor atoms of the 3-fpt^{2-} ligand (Table 2). The remaining water H atom along with the other two 3-fpt^{2-} O atoms are not involved in hydrogen bonding.




Figure 1
The coordination environment of the Cd^{II} atom in (I), showing the atom-labeling scheme. Displacement ellipsoids for non-H atoms are drawn at the 50% probability level. Hydrogen bonds are represented by dashed lines. [Symmetry codes: (i) $-x + 1, -y + 1, -z + 1$; (ii) $x - 1, y, z$; (iii) $-x + 1, -y + 2, -z + 1$.]




Figure 2
View of the crystal packing in (I). All H atoms, except for those of water, have been omitted for clarity. Hydrogen bonds are represented by dashed lines.

As is typical for metal phthalate-type complexes, the two carboxylate groups in (I) are not coplanar with the aromatic ring (Vaz *et al.*, 1996). One of the carboxylate groups makes a dihedral angle of $72.7(1)^\circ$, while the other is twisted at an angle of $31.5(1)^\circ$.

The 3-ft^{2-} ligand acts in a μ_4 -bridging fashion that links four symmetrically related Cd^{II} centers. The result is the formation of a 14-membered ring, an eight-membered ring, and a four-membered ring with $\text{Cd}\cdots\text{Cd}$ distances of $5.6292(4)$, $4.2213(3)$, and $3.7270(3)$ Å, respectively. The center of each ring corresponds to a crystallographic inversion center.

The aforementioned bridging of Cd^{II} centers in (I) results in the formation of thick polymeric sheets that are stacked along the c axis, with the benzene rings of the 3-ft^{2-} ligands projecting outward from each side (Fig. 2). Adjacent sheets are connected through π - π interactions involving the protruding aromatic rings of 3-ft^{2-} from successive layers, with a centroid-centroid distance of $3.67(2)$ Å between rings.

Experimental

All chemicals and solvents were purchased from commercial sources and used without further purification. 3-Fluorophthalic acid (3 mmol) was added to 100 ml of water and subsequently brought to pH 6.5 by the addition of 3 M NaOH with constant stirring. To this solution was added 10 ml of a 0.10 M solution of $\text{Cd}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O}$. Single crystals of (I) were obtained in two weeks after slow evaporation of this solution.

Crystal data

$[\text{Cd}(\text{C}_8\text{H}_3\text{FO}_4)(\text{H}_2\text{O})_2]$
 $M_r = 330.54$
Triclinic, $P\bar{1}$
 $a = 6.9460(5)$ Å
 $b = 7.2330(5)$ Å
 $c = 10.8199(8)$ Å
 $\alpha = 103.217(1)^\circ$
 $\beta = 93.351(1)^\circ$
 $\gamma = 115.066(1)^\circ$

$V = 471.86(6)$ Å³
 $Z = 2$
 $D_x = 2.326$ Mg m⁻³
Mo $K\alpha$ radiation
 $\mu = 2.34$ mm⁻¹
 $T = 294(2)$ K
Irregular cleavage fragment,
colorless
 $0.36 \times 0.22 \times 0.16$ mm

Data collection

Bruker SMART APEX CCD
diffractometer
 ω scans
Absorption correction: multi-scan
(SADABS; Bruker, 2003)
 $T_{\min} = 0.882$, $T_{\max} = 1.000$
(expected range = 0.607 – 0.688)

4370 measured reflections
1927 independent reflections
1893 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.012$
 $\theta_{\max} = 26.4^\circ$

Refinement

Refinement on F^2
 $R[F^2 > 2\sigma(F^2)] = 0.017$
 $wR(F^2) = 0.042$
 $S = 1.11$
1927 reflections
161 parameters
H atoms treated by a mixture of
independent and constrained
refinement

$w = 1/[\sigma^2(F_o^2) + (0.0169P)^2 + 0.4118P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.51$ e Å⁻³
 $\Delta\rho_{\min} = -0.40$ e Å⁻³

Table 1

Selected geometric parameters (Å, °).

$\text{Cd1}-\text{O2}^{\text{i}}$	2.3226 (15)	$\text{Cd1}-\text{O4}^{\text{iii}}$	2.2312 (16)
$\text{Cd1}-\text{O2}^{\text{ii}}$	2.3951 (15)	$\text{Cd1}-\text{O5}$	2.2866 (17)
$\text{Cd1}-\text{O3}$	2.2518 (16)	$\text{Cd1}-\text{O6}$	2.3308 (19)
$\text{O2}^{\text{i}}-\text{Cd1}-\text{O2}^{\text{ii}}$	75.64 (6)	$\text{O4}^{\text{iii}}-\text{Cd1}-\text{O3}$	115.46 (6)
$\text{O2}^{\text{i}}-\text{Cd1}-\text{O6}$	167.55 (6)	$\text{O4}^{\text{iii}}-\text{Cd1}-\text{O5}$	157.03 (6)
$\text{O3}-\text{Cd1}-\text{O2}^{\text{i}}$	109.30 (6)	$\text{O4}^{\text{iii}}-\text{Cd1}-\text{O6}$	98.11 (7)
$\text{O3}-\text{Cd1}-\text{O2}^{\text{ii}}$	161.87 (6)	$\text{O5}-\text{Cd1}-\text{O2}^{\text{i}}$	82.75 (6)
$\text{O3}-\text{Cd1}-\text{O5}$	85.61 (6)	$\text{O5}-\text{Cd1}-\text{O2}^{\text{ii}}$	77.62 (6)
$\text{O3}-\text{Cd1}-\text{O6}$	82.16 (7)	$\text{O5}-\text{Cd1}-\text{O6}$	93.58 (7)
$\text{O4}^{\text{iii}}-\text{Cd1}-\text{O2}^{\text{i}}$	81.61 (6)	$\text{O6}-\text{Cd1}-\text{O2}^{\text{ii}}$	91.96 (6)
$\text{O4}^{\text{iii}}-\text{Cd1}-\text{O2}^{\text{ii}}$	82.27 (6)		

Symmetry codes: (i) $-x + 1, -y + 1, -z + 1$; (ii) $x - 1, y, z$; (iii) $-x + 1, -y + 2, -z + 1$.

Table 2

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{O5}-\text{H5A}\cdots\text{O4}^{\text{iv}}$	0.765 (18)	2.076 (19)	2.840 (2)	175 (3)
$\text{O5}-\text{H5B}\cdots\text{O1}$	0.758 (19)	1.994 (19)	2.749 (2)	175 (4)
$\text{O6}-\text{H6B}\cdots\text{O1}^{\text{v}}$	0.764 (19)	2.040 (19)	2.804 (2)	179 (5)

Symmetry codes: (iv) $x - 1, y - 1, z$; (v) $x, y + 1, z$.

H atoms bonded to C atoms were placed in geometrically idealized positions and included as riding atoms [$C-H = 0.95 \text{ \AA}$ and $U_{iso}(H) = 1.2U_{eq}(C)$]. The water H atoms were located in difference maps and refined with the six O—H distances restrained to be equal with an effective s.u. of 0.015 \AA .

Data collection: *SMART-NT* (Bruker, 2003); cell refinement: *SAINT-Plus-NT* (Bruker, 2003); data reduction: *SAINT-Plus-NT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 1997); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997); molecular graphics: *SHELXTL* and *DIAMOND* (Brandenburg, 2005); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2000).

Financial support from the National Science Foundation (awards CHE-0314164 and CHE-0315152) is gratefully acknowledged.

References

- Brandenburg, K. (2005). *DIAMOND*. Version 3.0e. Crystal Impact GbR, Bonn, Germany.
- Bruker (2003). *SMART* (Version 5.625), *SAINT-Plus* (Version 6.45) and *SADABS* (Version 2.05). Bruker AXS Inc., Madison, Wisconsin, USA.
- Kim, Y. J., Suh, M. & Jung, D.-Y. (2004). *Inorg. Chem.* **43**, 245–250.
- Mehrotra, R. C. & Bohra, R. (1983). *Metal Carboxylates*. London: Academic Press.
- Rowsell, J. L. C., Millward, A. R., Kyo, S. P. & Omar, M. Y. (2004). *J. Am. Chem. Soc.* **126**, 5666–5667.
- Sheldrick, G. M. (1997). *SHELXS97* and *SHELXL97*. University of Göttingen, Germany.
- Sheldrick, G. M. (2000). *SHELXTL*. Version 6.1. Bruker AXS Inc., Madison, Wisconsin, USA.
- Vaz, J. L. L., Duc, G., Petit-Ramel, M., Faure, R. & Vittorei, O. (1996). *Can. J. Chem.* **74**, 359–364.
- Wasuke, M., Tomohiko, S., Tesushi, O., Chika, N. K. & Tohru, T. (2005). *J. Solid State Chem.* **178**, 2555–2573.
- Ye, B.-H., Tong, M.-L. & Chen, X.-M. (2005). *Coord. Chem. Rev.* **249**, 545–565.